

The National Animal Genetic Resources Center and Data Bank (NAGRC&DB)



"Animal genetic solutions for economic transformation"



BULL CATALOGUE

PREFACE

The National Animal Genetic Resources Centre and Data Bank (NAGRC&DB) is a body corporate under the Ministry of Agriculture, Animal Industry and Fisheries. It was established by an Act of Parliament in 2001, and is mandated to spearhead livestock breeding in the country. NAGRC&DB pursues scientifically based breeding with a goal to improve livestock genetics through discriminate crossbreeding, systematic selection of dams/sires, conservation of indigenous animal genetics, and generation and storage of breeding and production records. It also gives guidance on the breeds and the type of bulls to be used by farmers.

OUR VISION: A leader in profitable production and efficient delivery of animal genetic resources and services in Eastern Africa.

OUR MISSION: To establish a comprehensive and sustainable National Animal Breeding Program which meets the commercial and developmental interests of the actors along the livestock subsector value chains.

The bulls presented in this catalogue are bulls from which semen is collected and are housed at NAGRC&DB Bull Stud (a place where bulls are kept). They are exotic breeds that have been bred and selected under tropical conditions, specifically, in the East African countries. Exotic bulls bred under temperate conditions and transferred to tropical countries have frequently posed large problems due to lack of adaptation. Use of imported semen is slightly better than importing live animals. However, use of semen from bulls which have been raised in the country, the region, or in the tropical belt, is the best way to develop climate resilient stock. In other words, such an approach will enable us have animals which perform well (give more milk and beef) on our natural and improved pastures, and at the same time are more resistant to diseases and parasites.

This Bull Catalogue is a simplified version for farmers and stakeholders in the livestock value chain. The characteristics or traits presented can easily be understood to enable farmers make choices of bulls whose semen they can use on their farms. Extension workers, especially artificial insemination technicians, should accordingly assist farmers in making use of this catalogue.

Dr. Peter Beine Executive Director

GUIDANCE IN THE USE OF THE CATALOGUE

There are terms and graphics that have be used in describing the traits of the bulls which the reader needs to understand properly. The pedigree of the animals have been presented for both the dairy and beef bulls while the functional / linear descriptive traits have been presented for only dairy cattle.

PEDIGREE - means the recorded ancestry or lineage of an animal


PUREBRED – refers to the animal having a known ancestry. Bred from members of a recognized breed, strain, or kind without admixture of other blood over many generations

Sire: male parent of an animal

Dam: female parent of an animal

Grandsire: male grandparent of an animalGranddam: female grandparent of an animal

KLBO: Kenya Livestock Breeders Organization



FUNCTIONAL / LINEAR DESCRIPTIVE TRAITS

Functional traits are a species' morphological (form and structure), physiological (cell, tissue and organism functions), or phenological (seasonal biological events) characteristics which impact fitness indirectly via their effects on growth, reproduction and survival.

Each Linear descriptive trait rating is based on a measurement made by the classifier. In most cases, this is not an actual measurement made by a yardstick but by rating an animal's trait within a range of biological extremes. The traits are rated without regard to age, environment or stage of lactation. The 15 functional or Linear Descriptive traits that have been included in this catalogue are called primary traits. They have economic value and are practical traits to select for breed improvement. The illustrations at the end of the Catalogue will assist in explaining the individual trait being evaluated.

KEBEN Original name: KAP. KEBENEI LIABA 353RD Origin: Kenya Date of birth: 04.8.2015 KLBO Cert No. 23561. LXXVIII Breeder: Kapsoein Estates Ltd



Paternal Grand Sire (PGS)						
Conformation / Functional / Linear Descriptive Traits						
			-2	-1 0	1 2	
Stature	+1.45	Tall				
Strength	+1.52	Strong				
Body Depth	+1.54	Deep				
Dairy Form	+0.05	Open Rib				
Rump Angle	+1.54	Sloped				
Rump Width	+1.45	Wide				
Rear Legs-Side	+0.47	Sickle				
Rear Legs-Rear	-1.14	Hock In				
Foot Angle	-1.13	Low				
Feet & Legs Score	-0.78	Low				
F. Udder Attachment	-0.92	Loose				
Rear Udder Height	-0.49	Low				
Rear Udder Width	+0.24	Wide				
Udder Cleft	+0.94	Strong				
Udder Depth	-0.95	Deep				
Front Teat Placement	+0.42	Close				
Rear Teat P. Rear	+1.67	Close				
Teat Length	+1.41	Long				




Lactation No.	Days in Lactation	Total Yield (kg)
1.	301	6,350
2.	301	6,780
3.	305	7,128

COPPER Original name: Chemusian 16CC73 Origin: Kenya Date of birth: 24.04.16 KLBO Cert No. 23644. LXXVIII Breeder: Chemusian Co. Ltd



Sire							
Conformation / Functional / Linear Descriptive Traits							
			-2	-1	0	1 2	
Stature	-0.22	Short					
Strength	+0.80	Strong					
Body Depth	+1.05	Deep					
Dairy Form	+0.54	Open Rib					
Rump Angle	+0.51	Sloped					
Thurl Width	+0.77	Wide					
Rear Legs-Side	+0.50	Sickle					
Rear Legs-Rear	-0.56	Hock In					
Foot Angle	-0.61	Low					
Feet & Legs Score	-0.83	Low					
F. Udder Attachment	-0.81	Loose					
Rear Udder Height	+0.02	High					
Rear Udder Width	-0.16	Narrow					
Udder Cleft	-0.70	Weak					
Udder Depth	-1.71	Deep					
Front Teat Placement	-1.19	Wide					
Rear Teat P. Rear	-0.45	Wide					
Teat Length	+2.94	Long					




Lactation No.	Days in Lactation	Total Yield (kg)
1.	301	7,625
2.	301	7,428
3.	305	7,570

PUNGIN Original name: CHEMISIAN 16CC69 Origin: Kenya Date of birth: 02.04.2016 KLBO Cert No. 23638. LXXVIII Breeder: Chemusian Co. Ltd



Paternal Grand Sire (PGS)

Conformatio	n / Fur	ctional / I	inear	r Desc	riptiv	e Trai	ts	
			-2	-1	0		1	2
Stature	+0.64	Tall						
Strength	-0.56	Frail						
Body Depth	+0.25	Deep						
Dairy Form	+1.83	Open Rib						
Rump Angle	+0.65	Sloped						
Thurl Width	+0.52	Wide						
Rear Legs-Side	+0.57	Sickle						
Rear Legs-Rear	-1.76	Hock In						
Foot Angle	-0.45	Low						
Feet & Legs Score	-0.95	Low						
F. Udder Attachment	-0.86	Loose						
Rear Udder Height	-0.36	Low						
Rear Udder Width	+0.00	Wide						
Udder Cleft	+1.22	Strong						
Udder Depth	+0.49	Shallow						
Front Teat Placement	+0.13	Close						
Rear Teat P. Rear	+0.68	Close						
Teat Length	-0.46	Short						




Lactation No.	Days in Lactation	Total Yield (kg)
1.	301	7,625
2.	301	7,428
3.	305	7,570

TYSON Original name: GICHEHA TYSON 16GAMO15 ET Origin: Kenya Date of birth: 22.01.2017 KLBO Cert No. 23675. LXXVIII Breeder: Gicheha Farms Ltd



Parternal Grand Sire (PGS) Conformation / Functional / Linear Descriptive Traits +1.45 Tall Stature Strength +1.52 Strong +1.54 Deep Body Depth +0.05 Open Rib Dairy Form Rump Angle +1.54 Sloped Rump Width +1.45 Wide +0.47 Sickle Rear Legs-Side Rear Legs-Rear -1.14 Hock In Foot Angle -1.13 Low Feet & Legs Score -0.78 Low -0.92 Loose F. Udder Attachment Rear Udder Height -0.49 Low Rear Udder Width +0.24 Wide Udder Cleft +0.94 Strong Udder Depth -0.95 Deep Front Teat Placement +0.42 Close Rear Teat P. Rear +1.67 Close Teat Length +1.41 Long



Lactation No.	Days in Lactation	Total Yield (kg)
1.	315	6,810
2.	310	6,832
3.	305	7,035

SITBON


Original name: Chemusian 16CC32

Origin: KENYA | Date of birth: 12.02.2016 | KLBO Cert No. 23641. LXXVIII | Breeder: Chemusian Co.



Sire

Sir C								
Conformati	Conformation / Functional / Linear Descriptive Traits							
			-2	-1	()	1 2		
Stature	+0.64	Tall						
Strength	-0.56	Frail						
Body Depth	+0.25	Deep						
Dairy Form	+1.83	Open Rib						
Rump Angle	+0.65	Sloped						
Thurl Width	+0.52	Wide						
Rear Legs-Side	+0.57	Sickle						
Rear Legs-Rear	-1.76	Hock In						
Foot Angle	-0.45	Low						
Feet & Legs Score	-0.95	Low						
F. Udder Attachment	-0.86	Loose						
Rear Udder Height	-0.36	Low						
Rear Udder Width	+0.00	Wide						
Udder Cleft	+1.22	Strong						
Udder Depth	+0.49	Shallow						
Front Teat Placement	+0.13	Close						
Rear Teat P. Rear	+0.68	Close						
Teat Length	-0.46	Short						



Lactation No.	Days in Lactation	Total Yield (kg)
1.	287	5,969
2.	285	6,245
3.	275	6,038

AYRSHIRE


NIGEL JOYCE

Original name: Chemusian Joyce 4th

Origin: Kenya Date of birth: 03.02.2016 KLBO Cert No. 14697. LXXVIII Breeder: Chemusian Co. Ltd



Sire								
Conformation / Functional / Linear Descriptive Traits								
			-2	-1	()		1	2
Stature	+1.30	Tall						
Strength	-0.10	Frail						
Body Depth	+0.90	Deep						
Dairy Form	+1.30	Open Rib						
Rump Angle	+0.40	Sloped						
Rump Width	+1.20	Wide						
Rear Legs-Side	+0.70	Sickle						
Foot Angle	-0.70	Low						
F. Udder Attachment	+0.20	Strong						
Rear Udder Height	+0.80	High						
Rear Udder Width	+0.30	Wide						
Udder Cleft	+0.70	Strong						
Udder Depth	+0.00	Shallow						
Front Teat Placement	+1.20	Close						
Teat Length	-0.40	Short						




Lactation No.	Days in Lactation	Total Yield (kg)
1.	290	6,710
2.	270	6,576
3.	275	6,739

AYRSHIRE

CHEMUS Original name: Chemusian Mazalan Origin: Kenya Date of birth: 22.02.2016 KLBO Cert No. 14703. LXXVIII Breeder: Chemusian Co. Ltd



Sire								
Conformation / Functional / Linear Descriptive Traits								
			-2	-1	0		1	2
Stature	-0.30	Short						
Strength	+0.00	Strong						
Body Depth	+0.00	Deep						
Dairy Form	+0.60	Open Rib						
Rump Angle	+1.20	Sloped						
Thurl Width	-0.60	Narrow						
Rear Legs-Side	-0.20	Posty						
Foot Angle	+0.70	Steep						
F. Udder Attachment	-1.60	Loose						
Rear Udder Height	-0.20	Low						
Rear Udder Width	+0.00	Wide						
Udder Cleft	+0.70	Strong						
Udder Depth	-1.10	Deep						
Front Teat Placement	+1.10	Close						
Teat Length	-0.50	Short						




Lactation No.	Days in Lactation	Total Yield (kg)
1.	290	5,487
2.	270	6,340
3.	278	6,723

AYRSHIRE

KEFA Original name: Chemusian Mazalan Origin: Kenya Date of birth: 13.01.2017 KLBO Cert No. 14693. LXXVIII Breeder: Gicheha Farms Ltd



Sire								
Conformat	Conformation / Functional / Linear Descriptive Traits							
			-2	-1		()	1	2
Stature	-0.30	Short						
Strength	+0.10	Strong						
Body Depth	+0.10	Deep						
Dairy Form	+0.70	Open Rib						
Rump Angle	+1.30	Sloped						
Thurl Width	-0.60	Narrow						
Rear Legs-Side	-0.10	Posty						
Foot Angle	+0.70	Steep						
F. Udder Attachment	-1.60	Loose						
Rear Udder Height	-0.30	Low						
Rear Udder Width	+0.10	Wide						
Udder Cleft	+0.60	Strong						
Udder Depth	-1.20	Deep						
Front Teat Placement	+1.10	Close						
Teat Length	-0.40	Short						



Lactation No.	Days in Lactation	Total Yield (kg)
1.	310	5,526
2.	305	7,566
3.	308	8,200

JERSEY


FOUNTAIN

Original name: KIGWA JUBILEE 14/01

Origin: Kenya | Date of birth: 01.10.2014 | KLBO Cert No. 10391. LXXVIII | Breeder: Kigwa Estate



Sire Conformation / Functional / Linear Descriptive Traits Type %R 64 D / H Stature -0.64 Short -0.26 Capacity Low Rump Angle +0.05 Sloped Rump Width -0.25 Narrow +0.08Curved Legs Udder Support +0.10 Strong Fore Udder +0.11 Strong Rear Udder +0.31Strong Front Teat Placement Close +0.17Rear Teat Placement +0.13Close Udder Overall +0.29 Desirable Dairy Conformation -0.25 Undesirable




Lactation No.	Days in Lactation	Total Yield (kg)
1.	300	5,250
2.	297	5,438
3.	302	5,688

JERSEY

GROOVE Original name: Avalon Origin: Kenya Date of birth: 07.08.2015 KLBO Cert No. 10303. LXXVII Breeder: Avalon Jerseys



Sire							
Conformation	Conformation / Functional / Linear Descriptive Traits						
			-1	()	1	
Stature	-0.64	Short					
Capacity	-0.26	Low					
Rump Angle	+0.05	Sloped					
Rump Width	-0.25	Narrow					
Legs	+0.08	Curved					
Udder Support	+0.10	Strong					
Fore Udder	+0.11	Strong					
Rear Udder	+0.31	Strong					
Front Teat Placement	+0.17	Close					
Rear Teat Placement	+0.13	Close					
Udder Overall	+0.29	Desirable					
Dairy Conformation	-0.25	Undesirable					



Lactation No.	Days in Lactation	Total Yield (kg)
1.	295	5,290
2.	300	5,464
3.	302	5,650

GUERNSEY

BRIGHT Original name: GENERAL BRIGHT Origin: Uganda Date of birth: 12.03.2015 ID No.: 326 Breeder: NAGRC&DB




Sire **Conformation / Functional / Linear Descriptive Traits** PTAT +0.00 86%R UDC+0.1 FLC-1.7 108 D / 44 H +0.30 Tall Stature Strength -1.10 Frail Body Depth -0.30 Shallow +0.70 Open Rib Dairy Form Rump Angle +0.00 Sloped -2.20 Narrow Thurl Width Rear Legs-Side +0.10 Sickle Rear Legs-Rear -0.10 Hock In Foot Angle -1.10 Low F. Udder Attachment +0.20 Strong Rear Udder Height +1.20 High Rear Udder Width -0.40 Narrow Udder Cleft +0.10 Strong Udder Depth +1.30 Shallow Front Teat Placement -2.00 Wide +3.10 Teat Length Long



Lactation No.	Days in Lactation	Total Yield (kg)
1.	297	6,050
2.	300	7,020
3.	303	7,160


YOANI 2502					
Original name: 2502					
Origin: Kenya	Date of birth: 07.02.2018	KLBO Cert No. 2833.LXXXII	Breeder: Stanley & Son Ltd.		






YOANI 2517					
Original name: YOANI BIY 2517					
Origin: Kenya Date of birth: 17.04.2018 KLBO Cert No. 2832.LXXXII Breeder: Stanley & Son Ltd.					






YOANI 2564 Original name: YOANI BIY 2564 Origin: Kenya Date of birth: 26.11.2018 KLBO Cert No. 2935.LXXXII Breeder: Stanley & Son Ltd.






YOANI 2613						
	Original name: YOANI BIY 2613					
Origin: Kenya	Origin: Kenya Date of birth: 27.06.2019 KLBO Cert No. 2931.LXXXIII Breeder: Stanley & Son Ltd.					






DELO 258					
Original name: 258					
Origin: Kenya Date of birth: 23.08.2019 KLBO Cert No. PB/2344 Breeder: DELORAINE Estates					






DELO 261						
Original name: 261						
Origin: Kenya						






KARAMA 9084				
Original name: 9084				
Origin: Kenya	Date of birth: 27.05.2019	KLBA Cert No. PB/2383	Breeder: EL KARAMA Sahiwals Ltd,	





KARAMA 9091				
Original name: 9091				
Origin: Kenya	Date of birth: 05.06.2019	KLBA Cert No. PB/2375	Breeder: EL KARAMA Sahiwals Ltd.	






FUNCTIONAL / LINEAR DESCRIPTIVE TRAITS

Functional traits are a species' morphological (form and structure), physiological (cell, tissue and organism functions), or phenological (seasonal biological events) characteristics which impact fitness indirectly via their effects on growth, reproduction and survival.

Each Linear descriptive trait rating is based on a measurement made by the classifier. In most cases, this is not an actual measurement made by a yardstick but by rating an animal's trait within a range of biological extremes. The traits are rated without regard to age, environment or stage of lactation. The 15 functional or Linear Descriptive traits that have been included in this catalogue are called primary traits. They have economic value and are practical traits to select for breed improvement.

PARTS OF A COW



DAIRY CHARACTER

DAIRY FORM

IMPORTANCE:

Shows the cow's milking ability. Under good feeding conditions, less excess flesh and fat is an indication of a cow's ability to convert feeds more into milk than into beef. Sharpness and flatness of bone, openness and slant of rib, and length of neck are evaluated by the classifier to assign a dairy form rating

NOT DESIRED



EXTREMELY TIGHT

- Tighter, closer ribbed
- Short thick fleshy neck
- Excess flesh and fat especially in the throat, brisket or dewlap
- · Lacks angularity

INTERMEDIATE FORM

DESIRED



EXTREMELY OPEN

- · More openness of ribs
- · Long lean neck
- Less excess flesh and fat
- Very angular


FERTILITY

RUMP ANGLE

IMPORTANCE:

Indicates how good the drainage of the reproductive tract can be. Ideal rump angle is a slight slope (1.2 inches below the tip). Observing the animal from the side, the classifier notes the angle of the rump structure from hook (hips) to pins. A rating is given based on the degree to which the pins are higher or lower than the hooks.

NOT DESIRED



HIP LOWER THAN PIN

DESIRED MODERATE RUMP ANGLE



SLIGHT SLOPE FROM HIP TO PIN

RELATIVELY DESIRED HIGH RUMP ANGLE




HIP HIGHER THAN PIN

- Extreme slope from the hip to the pin
- Not desirable for the height of the rear udder attachment

RUMP WIDTH

IMPORTANCE: Indicates calving ease. The wider the rump, the easier it is for the cow or heifer to calve down (deliver a calf). It is determined through evaluating the distance between the inside wall of pins.


NOT DESIRED



EXTREMELY NARROW

INTERMEDIATE

DESIRED



EXTREMELY WIDE


BODY CAPACITY (ROUGHAGE CONSUMPTION)

BODY DEPTH

IMPORTANCE: Indicates the cow's capacity to consume large amounts of feeds leading to more milk production. A wide and big mouth, and deep flanks also promote more feed intake. The classifier evaluates depth of


body by looking primarily at the rib cage.

NOT DESIRED




EXTREMELY SHALLOW

- Barrel: long, deep, and wide
- Long and wide ribs



INTERMEDIATE

DESIRED



EXTREMELY DEEP BODY

BODY CAPACITY (STRENGTH)

CHEST WIDTH

IMPORTANCE: Indicates the cow's ability to sustain high production and good general health. In addition to width of the chest, muzzle width and substance of bone in the cow's front end comprise strength.


NOT DESIRED



NARROW & FRAIL

INTERMEDIATE

DESIRED




WIDE & STRONG

LONGEVITY

REAR LEGS SET (SIDE VIEW)


IMPORTANCE: Indicates the durability of the legs and feet hence the longevity and efficiency of a cow. The cow should have good mobility and be able to move smoothly. The classifier evaluates rear legs from the side, noting the amount of set to the hock joint.

NOT DESIRED




POSTY AND STRAIGHT Causes too much stress on the legs leading to aggravation of joints

DESIRED



MODERATE SET (CURVE) TO THE HOCK

NOT DESIRED



SICKLECauses too much stress on the leg muscles and tendons

FOOT ANGLE

IMPORTANCE: Indicates the cow's durability and mobility. It also determines how frequent a cow's feet needs trimming. Foot angle ratings are based on the steepness of the angle of the foot as viewed from the side.


NOT DESIRED



LOW ANGLED Shallow heel

INTERMEDIATE

DESIRED




VERY STEEP Steep claw with a deep heel

MILK PRODUCTION (UDDER)

REAR UDDER HEIGHT

IMPORTANCE: Indicates the capacity of the udder. The higher the attachment of the udder, the more udder capacity hence more milk production. The classifier assigns a rating to rear udder height by viewing the cow from the rear. The point of rear udder attachment determines rear udder height.

NOT DESIRED



VERY LOW ATTACHED UDDER Low udder capacity



INTERMEDIATE

DESIRED



HIGHLY ATTACHED UDDER More capacious udder

UDDER CLEFT – CENTRAL LIGAMENT

IMPORTANCE: Indicates how easy a cow can be milked and also its susceptibility to udder injury. A clearly defined halving in the udder makes milking easy. A stronger center support minimizes udder injury. A deep udder cleft is an indicator of a strong median suspensory ligament. The strong udder support prevents the udder from becoming too deep. Also helps in proper teat placement. Udder cleft is evaluated by viewing the bottom of the udder.

NOT DESIRED



WEAK CLEFT

- Convex to flat floor
- Udder lacking cleavage



INTERMEDIATE Slight definition

DESIRED



STRONG CLEFT

- Strong median (central) suspensory ligament support
- Deep defined halving

MILK PRODUCTION (UDDER)

FORE UDDER ATTACHMENT

IMPORTANCE:

Important to herd life as it affects udder depth and susceptibility of the udder to injury. To determine fore udder attachment ratings, the classifier looks at the strength of the attachment to the body wall by lateral ligaments.

NOT DESIRED



WEAK AND LOOSE

DESIRED



INTERMEDIATE

RELATIVELY DESIRED



EXTREMELY STRONG AND TIGHT


- Udder held tighter to the body wall
- Very good for longevity, but negatively correlated to milk production

MILK PRODUCTION (UDDER)

UDDER DEPTH


IMPORTANCE: Indicates the capacity of the udder and susceptibility of the udder to injury and mastitis Cows with moderate udder depth stay in the herd longer.

NOT DESIRED




DEEP UDDER FLOOR BELOW HOCKS Teats susceptible to injury

DESIRED



UDDER FLOOR ABOVE HOCKS

DESIRED




SHALLOW – EXTREME HEIGHT OF UDDER FLOOR ABOVE HOCKS

MILK PRODUCTION (TEATS)

FRONT TEAT PLACEMENT


IMPORTANCE: Proper teat placement makes milking easy and milking completely reduces susceptibility of the teat to injury/mastitis. The rating is assigned as the classifier views the teats from the rear.

NOT DESIRED




EXTREME PLACEMENT ON OUTSIDE OF QUARTER
Teats placed on the edge of the quarters

DESIRED



CENTRALLY PLACED ON QUARTER

NOT DESIRED




EXTREME PLACEMENT INSIDE QUARTERTeats hang too close together

REAR TEAT PLACEMENT


IMPORTANCE: Proper teat placement makes milking easy and reduces susceptibility of the teat to injury

NOT DESIRED




EXTREME PLACEMENT ON OUTSIDE OF QUARTERTeats placed on the edge of the quarters

DESIRED



CENTRALLY PLACED ON QUARTER

NOT DESIRED




EXTREME PLACEMENT INSIDE QUARTERTeats hang too close together

MILK PRODUCTION (TEATS)

TEAT SIZE AND SHAPE


IMPORTANCE: Proper teat placement makes milking easy and reduces susceptibility of the teat to injury. Unusual size and shape result in milking problems.

NOT DESIRED




SHORT Short teats make milking difficult

DESIRED



INTERMEDIATE

NOT DESIRED



LONGLong teats are susceptible to injury


STATURE

IMPORTANCE:

Related to udder depth. The taller the cow, the higher the udder floor above the hocks (higher udder depth) hence less chances of injuries. Taller cows tend to be bigger, feed more and may be produce more.

- Adequate height including length in leg bones with a long bone pattern is desirable
- Height at withers and hips should be relatively proportionate

Short: 1.30 m Intermediate: 1.42 m Tall: 1.54 m





Measured from the top at a point in the middle of the backbone (spine) between hip bone and the vertebrae to the ground.

PARTS OF A COW

PARTS	DESCRIPTION AND DESIRABLE CHARACTERISTICS
Back	Comprises of the chine and the loin. A cow's back should be straight and strong.
Barrel	The entire area below the cow's back, from withers to hips. The barrel should be long, deep, and wide.
Bridge of nose	The part of the head that begins just below the eyes and continues all the way to the muzzle. The bridge of nose should be straight.
Brisket	Part of the cow's chest, the area between, and slightly in front of the legs. Along with the dewlap, the brisket should be lean and clean cut.
Chest Floor	The area between and directly behind the front legs. The chest should be deep and wide.
Chine	The chine relates to the upper spine of a human. It should be sharp and prominent
Crops	The area directly behind the top of the shoulder blade. The crops should have adequate fullness.
Dewclaw	The two small, hoof-like points on the back side of each leg just above each pastern.
Dewlap	The loose skin along the lower part of the neck. The dewlap should be lean and clean cut.
Flank	The area between the rear ribs and the rear legs that adds depth to the body. The flank should be deep and refined.
Fore Udder	The two front quarters of the udder. The fore udder should be firmly attached with moderate length and ample capacity.
Fore Udder Attachment	The line along the top of the fore udder where the fore udder attachment should blend smoothly into the body wall and have adequate length.
Forehead	The area between the eyes. A cow's forehead should be broad and moderately dishes
Heart Girth	The diameter of the front end of the barrel. Almost called the chest. A cow s heart girth should be deep and have a wide floor with well-sprung fore ribs blending into the shoulders.
Heel	Lies at the back of the hoof. A cow's heel should be deep.

Hip Large, bony structures that stick out on each side of the cow.

Sometimes called hooks, or hip bones. The hips should be wide and

prominent.

Hock A joint, which corresponds to your ankle and only bends forward.

Hocks should be cleanly molded, free from coarseness and puffiness

with adequate flexibility.

Hoof The hoof of a cow has toes, a split toe. The hoof should be well

rounded with closed toes.

Identify this characteristic by comparing them almost exactly to the Jaw

same part of your head and neck. A cow s jaw should be strong and

wide.

Knee The cow's kneel is most similar to our wrist. Kneels should be straight

and free from swelling and coarseness.

Loin Found between the chine and the rump. The loin should be broad,

strong, and nearly level.

Milk Veins Lie along the underbody of the cow, prior to the fore udder.

Milk Wells Lie along the underbody of a cow, along the chest floor.

Muzzle The part of the head that looks like it ought to be the cow's nose. A

cow's muzzle should be strong and have large, open nostrills.

Neck Identify this characteristic by comparing them almost exactly to the

same part of your head and neck. The neck needs to be long, lean, and

smoothly blended into the shoulders.

Located between the dewclaws and the top of the hoof, the pastern **Pastern**

works like a shock absorber with every step the cow takes. Pasterns

should be short and strong with some flexibility.

Pin Bones Sometimes called pins, are similar to our hips socket. The pins should

be clearly difined, wide apart, and set slightly lower than the hip bones.

Point of Elbow The front leg joint located above the cow's knee. This corresponds

exactly to our elbow. The point of elbow needs to be set firmly against

the body wall.

Point of Just like our shoulder, the cow s point of shoulder blade. The point of Shoulder

shoulder should be clearly difined with the entire shoulder blade set

tightly against the body wall.

Poll The part of the cow's head located between the horns.

Real Udder The two rear quarters of the udder. The rear udder should be wide and

high while being firmly attached with uniform width from top bottom and

slightly rounded to the udder floor.

Rear Udder Attachment

The point between the rear legs and the thighs where the real udder is secured is the real udder attachment. The real udder attachment should be wide, high, and have adequate symmetry and fulness at the top.

Ribs The cow's ribs create a barrel shape in the middle part of the cow. Her

ribs should be wide apart, flat, deep, and slanted towards the rear.

Rump Begins at the cows hips and ends at the pin bones. Serves as the

starting point for the leg and the enclosure for the cows reproductive system. The rump should be long and wide throughout with the pin

bones slightly lower than the hip bones.

Sole Lies along the bottom of the hoof. The sole should be flat.

Stifle Halfway between the hock and the thurl, it is the cows kneel joint and

only bends backward.

Swich The long, bushy hair at the end of the tail.

Tail It should be free from coarseness.

Tailhead Acts as the starting point for the cows tail. The tailhead should be set

slightly above and neatly between a cow s pins.

Teats Four, located at the base of the udder allow milk to be removed from

each quarter. Teats should have cylindrical shape and be of uniform

size with medium length and diameter.

Thigh The area long the upper part of the back side of the rear legs. Thighs

are lean, incurving to flat, and wide apart from the rear.

Throat Identify this characteristic by comparing them almost exactly to the

same part of your head and neck. The throat should be clean-cut.

Thurl A joint which attaches the rear leg to the rump. The thurls need to be

wide apart and centrally placed between the hip bones and the pin

bones.

Top line Every part of the cow that you might want to call the back combines to

form the topline, from the withers to the pin bones. The topline should

be straight and strong.

Withers The area where the chine begins. The withers should be sharp.



For more information Please contact:

National Animal Genetic Resources Centre & Data Bank Plot 98-106 Nsamizi Road, Entebbe

P.O. Box 183, Entebbe-Uganda

: +256 414 320 831
 : info@nagrc.go.ug
 : www.nagrc.go.ug
 : @NAGRC_DB